

# Sonoelastography in Nonalcoholic Fatty Liver Disease

Arnabjyoti Deva Sarma<sup>1\*</sup>, Jibon Sharma<sup>2</sup>, Moitrayee Devi<sup>3</sup>

## ABSTRACT

**Background of the Study:** It is an imaging technology sensitive to tissue stiffness that was first described in the 1990s. Nonalcoholic fatty liver disease (NAFLD) is increasingly common in Western nations of the world. It is the most common form of chronic liver disease, affecting about one-quarter of the population, in the United States. **Materials and Methods:** Prospective studies of 100 patients with clinical impression of alcoholic were examined by Philips ultrasound imaging. **Results:** A total of 100 patients satisfying the inclusion criteria were included in the study. The age range of patients was from 18 years to elderly with male predominance, male 64 (64%), and female 36 (36%). **Conclusion:** In this study, a non-invasive and cost-effective technique, without any radiation hazards, to measure the elastic property of liver tissue using ultrasound imaging modality called as sonoelastography in Indian population for assessment of disease severity and grading of NAFLD.

**Keywords:** Biopsy, Elastography, Fibrosis, Liver parenchyma, Ultrasound

*Asian Pac. J. Health Sci.*, (2022); DOI: 10.21276/apjhs.2022.9.4.20

## INTRODUCTION

Fatty liver disease is emerging as an important cause of liver disease in Northern India. Epidemiological studies suggest prevalence of non-alcoholic fatty liver disease (NAFLD) in around 9–32% of general population in India with higher prevalence in those with overweight/obesity and diabetes/pre-diabetes.<sup>[1]</sup> NAFLD is a common clinicopathological condition characterized by significant lipid deposition in the hepatocytes of the liver parenchyma. Ludwig *et al.*<sup>[2]</sup> in 1980 originally coined the term non-alcoholic steatohepatitis, "NASH" to describe the histologic evidence suggestive of alcoholic hepatitis on liver biopsy (i.e., steatosis and lobular inflammation) but no history of alcohol abuse. Many of these patients were females (60%) and the majorities were obese (90%). Several other terms have been used to refer to this entity, including pseudoalcoholic liver disease, alcohol-like hepatitis, diabetic hepatitis, non-alcoholic Laennec's disease, and steatonecrosis. However, seeing that the disease represents a spectrum of pathology, the umbrella term "NAFLD," first introduced in 1986, became the preferred one.<sup>[3]</sup>

The increased prevalence of fatty liver disease relates directly to the obesity epidemic seen in these populations. The most of the patients who come to medical attention with NAFLD are identified as a result of incidentally discovered elevated liver enzymes; alanine transaminase (ALT), aspartate transaminase (AST), and/or fatty liver on ultrasound examination. When patients are symptomatic, symptoms include fatigue or a vague right upper quadrant discomfort. Aminotransferases are only mildly (1.5–2 times the upper limit of normal) elevated, ALT is generally higher than AST. The most of the studies have shown that many patients can have advanced fibrosis with NASH and even cirrhosis due to NASH with normal liver enzymes. This indicates that the prevalence of the disease is likely to be even greater than previously suspected.<sup>[4]</sup>

The degree of liver fibrosis estimation helps to determine the prognosis and optimal treatment for fatty liver disease. Liver biopsy is taken as the standard reference for the diagnosis and staging of fibrosis and is considered the gold standard in diagnosis. This is the only reliable tool for distinguishing NASH from simple steatosis and for grading and staging the disease, providing important information about severity of steatosis, lobular inflammation, hepatocellular ballooning, and degree of fibrosis.<sup>[5]</sup>

<sup>1</sup>Department of Radiology, Assam Downtown University, Guwahati, Assam, India

<sup>2</sup>Department of Oncology, State Cancer Institute, Gauhati Medical College & Hospital, Guwahati, Assam, India

<sup>3</sup>Department of Microbiology, Assam Downtown University, Guwahati, Assam, India

**Corresponding Author:** Arnabjyoti Deva Sarma, Department of Radiology, Assam Downtown University, Guwahati, Assam, India. E-mail: sarma.arnab1990@gmail.com

**How to cite this article:** Sarma AD, Sharma J, Devi M. Sonoelastography in Nonalcoholic Fatty Liver Disease. *Asian Pac. J. Health Sci.*, 2022;9(4):95-98.

**Source of support:** Nil

**Conflicts of interest:** None.

**Received:** 02/02/2022 **Revised:** 17/03/2022 **Accepted:** 17/04/2022

However, this procedure is invasive and associated with high risk of complications and significant sampling and observer error because of the evaluation of a very small volume of tissue sample which is only 1/50000 of the total volume, thus limiting the accuracy for determination of fibrosis stage of liver. Liver biopsy is also unpleasant for patients and usually requires hospitalization and for the adequate interpretation of the specimen requires a pathologist with expertise in hepatopathology, which altogether makes it a costly and time-consuming procedure. In a well-designed study by Ratzu *et al.*,<sup>[6]</sup> the negative predictive value of a single biopsy for the diagnosis of NASH was calculated to be only 74%.

Hence, several non-invasive methods of histological assessment including clinical, biochemical, and radiological methods have been developed. Baršić *et al.*<sup>[7]</sup> worked on the overview and developments in non-invasive diagnosis of non-alcoholic fatty liver disease and concluded that even though, the gold standard still used in almost all of the studies is liver biopsy, but the question that remains is whether we are actually able to accurately assess the performance of non-invasive methods when the gold standard by itself has significant flaws. The study concludes of that there is a need for a new much efficient and a non-invasive tool for the diagnosis of NAFLD.

In this study, I have discussed on a non-invasive and cost-effective technique, without any radiation hazards, to measure the

elastic property of liver tissue using ultrasound imaging modality called as sonoelastography in Indian population for assessment of disease severity and grading of NAFLD.

## MATERIALS AND METHODS

The common methods for sonoelastography examination are as follows:

- The diagnosis of NAFLD is based on the clinical findings and routine ultrasound examination of the upper abdomen.
- The healthy controls can be selected based on the criteria; age older than 18 years, no H/O alcohol intake, no known H/O diabetes, hypertension, hypothyroidism, Jaundice in the past, body mass index (BMI) <25 kg/m<sup>2</sup>, normal liver function tests (LFT), normal lipid profile, negative viral markers for Hepatitis B (HepB) and Hepatitis C (HepC), and normal liver on routine USG examination.
- The patients of NAFLD can select based on the criteria; age older than 18 years, no H/O alcohol intake, no known H/O diabetes, hypertension, hypothyroidism, jaundice in the past, negative viral markers for HepB and HepC, and fatty liver on routine USG examination.
- Clinical profile of all the patients including BMI must be noted, liver function tests and lipid profile can be done in all patients. Grading of fatty liver was done based on the USG findings.
- Sonoelastography can be performed on all the healthy controls and the patients and their values must be noted.
- Liver biopsy can be done as and when indication clinically.

## Technique of Sonoelastography

### *Preparation of the patients*

Before starting the scan, procedure was explained to the patient in his/her vernacular language to allay any kind of fear and anxiety. A written consent about the explained details of investigation was taken.

### *Scanning procedure*

Sonoelastography was performed on ultrasound. Shear wave elastography (SWE) of liver can be done using convex probe after making proper position of the patient and appropriate positioning of the US probe. Eight acquisitions can be performed by placing the SWE box at different areas of liver. The readings of liver stiffness and velocity must be noted in kilopascals (kPa) and metres per second (m/s), respectively. A median value may then be determined and used as a representative measurement of the liver elasticity. Findings of shear wave sonoelastography of healthy controls can be considered as normal values for the purpose of comparison with the values obtained in patients of NAFLD. The findings of SWE will then be correlated with clinical profile and laboratory investigations and with liver biopsy wherever done.

## RESULTS

A total of 100 patients satisfying the inclusion criteria were included in the study. The age range of patients was from 18 years to elderly with male predominance, male 64 (64%), and female 36 (36%). Non-alcoholic fatty liver was the most common clinical diagnosis constituting (80%) cases.

## DISCUSSION

Imaging modalities frequently used in the diagnosis of NAFLD include ultrasound, computed tomography, and magnetic resonance imaging (MRI). While they are all very sensitive (80–100%) and specific in detection of steatosis, none of them can effectively distinguish simple steatosis from NASH or determine the degree of fibrosis.<sup>[8]</sup> Ever since hippocrates, physicians have used palpation to detect differences in tissue stiffness as an aid to diagnosis. Elastography depends on the same differences in mechanical properties between healthy and abnormal tissues but uses imaging to detect these differences at depths not reachable by manual palpation. Elastography techniques have been developed for both ultrasound and MRI. A major surge in development of ultrasonic methods and devices for elasticity imaging started in the late 1980s and early 1990s.<sup>[9–12]</sup> Beginning in the late 1990s, several laboratories developed an alternative approach for remote probing tissue elasticity using acoustic radiation force.<sup>[13–15]</sup> Acoustic radiation force is the time-average force exerted on an object by an acoustic wave.

Until recently, elasticity imaging was largely a research method used by a few select institutions having the special equipment needed to perform the studies. Since 2005, however, increasing numbers of mainstream manufacturers have added elastography to their ultrasound systems so that today, the majority of manufacturers offer some sort of elastography or tissue stiffness imaging on their clinical systems. Now, it is safe to say that elasticity imaging may be performed on virtually all types of focal and diffuse diseases. Transient elastography (TE) uses an external actuator to provide a single cycle of low-frequency (typically around 50 Hz) vibration and ultrasound methods to track the resulting motion. This type of excitation generates four types of waves including a compression and shear wave in the medium. The compression and shear waves are separated in time because the longitudinal wave speed is, in most cases, much faster than the shear wave. Applying a sinusoidal excitation at low frequencies with a mechanical actuator with a small cylindrical footprint can cause biases due to wave diffraction from the cylindrical source. Using a transient excitation avoids these biases so that the shear wave can be separated from the compression wave and any reflected waves.<sup>[16,17]</sup> TE has been used both as a one-dimensional measurement technique<sup>[17,18]</sup> as well as a two-dimensional imaging technique.<sup>[19]</sup> The technique has been developed for measurement of stiffness in the liver in a product called Fibro Scan<sup>®</sup> manufactured by EchoSens.<sup>[20]</sup> TE has also been employed for measuring stiffness in phantoms, skeletal muscle, breast, skin, and blood clots. Shear wave elasticity imaging (SWEI) was initially described theoretically by Sarvazyan *et al.*<sup>[13]</sup> and investigated experimentally by Nightingale *et al.*<sup>[21]</sup> and their combined work was a catalyst for many investigations and development of other techniques that followed. The use of modulated ultrasound was proposed to produce an acoustic radiation force imaging (ARFI) that would create shear waves that could be detected by optical, acoustic, ultrasound, or MRI methods. The radiation force acts as a “virtual finger” that may be used to palpate the organ from the inside, thereby replacing the physician’s fingers on the surface of the body or organ.

In SWEI and ARFI imaging, the radiation force is focused at a single location. An extension of these methods is to focus the radiation force in one location and then change the depth of the

**Table 1** : Distribution of abnormalities in various age groups

<i>USG Diagnosed</i>	18–30 years	31–45 years	45–60 years	>60 years	Total
Non-alcoholic fatty liver	7	9	0	0	16
	10	4	1	1	17
	0	0	1	0	1
	0	0	0	0	2
	1	5	4	0	10
	0	0	3	0	3
	4	2	2	1	15

focal location so that the shear waves created from multiple focal locations constructively interfere to make a conical shear wave.<sup>[14]</sup> This method is called the supersonic shear imaging because the radiation force focal point moves at a rate that is faster than the speed of the shear wave in the medium, providing credence to the “supersonic” nomenclature. The shear wave created forms a kind of Mach cone and the Mach number of the excitation can be adjusted make the shear wave directionally oriented. Detection is accomplished by ultrafast real-time ultrasound imaging, and the data are displayed in kPa and velocity in m/s. This method has the advantage of being quantitative, reproducible, and operator independent. It is, therefore, suitable for monitoring changes over time. Nevertheless, MRI is more sensitive than ultrasound in detecting lesser degrees of hepatic steatosis, and new techniques in MRI are constantly being developed that provide additional data on different tissue parameters. One of them is magnetic resonance (MR) elastography, which estimates liver tissue stiffness. This technique was shown to have an excellent predictive value for excluding fibrosis, while sensitivity and specificity for discriminating between mild and more severe fibrosis were around 85%.<sup>[22,23]</sup> The future advances in MRI technology including hepatic flow parameters and diffusion-weighted MRI may hopefully provide more MR-based tools for liver fibrosis detection. However, MRE may not be widely available for evaluating liver stiffness and its use as a screening tool may be time consuming and cost-prohibitive compared to ultrasound-based techniques.

**CONCLUSION**

On the basis of findings, we can conclude that:

- A standard technique of liver sonoelastography can be established.
- Can determine the normal values of liver sonoelastography in our healthy controls.
- Can easily evaluate the role of sonoelastography of liver in patients of NAFLD.
- Easily compare the sonoelastography findings with the clinical profile, BMI, LFT, lipid profile, and wherever required with liver biopsy

In this study, a non-invasive and cost-effective technique, without any radiation hazards, called as sonoelastography in Indian population for assessment of disease severity and grading of NAFLD can easily determine compared to other medical devices.

**REFERENCES**

1. Duseja A. Nonalcoholic fatty liver disease in India: A lot done, yet more required! *Indian J Gastroenterol* 2010;29:217-25.
2. Ludwig J, Viggiano TR, McGill DB, Oh BJ. Nonalcoholic steato-hepatitis: Mayo Clinic experiences with a hitherto unnamed disease. *Mayo Clin*

- Proc 1980;55:434-8.
3. Sheth SG, Gordon FD, Chopra S. Nonalcoholic steatohepatitis. *Ann Intern Med* 1997;126:137-45.
4. Eckel RH. The metabolic syndrome. In: *Harrison's Principles of Internal Medicine* 18<sup>th</sup> ed., Ch. 242; 2018. p. 1992-1997.
5. Neuschwander-Tetri BA, Clark JM, Bass NM, Van Natta ML, Unalp-Arida A, Tonascia J, Zein CO, *et al.* Clinical, Laboratory and histological associations in adults with nonalcoholic fatty liver disease. *Hepatology* 2010;52:913-24.
6. Ratziu V, Charlotte F, Heurtier A, Gombert S, Giral P, Bruckert E, *et al.* Sampling variability of liver biopsy in nonalcoholic fatty liver disease. *Gastroenterology* 2005;128:1898-906.
7. Baršić N, Ivan L, Lea SD, Vedran T, Marko D. Overview and developments in noninvasive diagnosis of nonalcoholic fatty liver disease. *World J Gastroenterol* 2012;18:3945-54.
8. Saadeh S, Younossi ZM, Remer EM, Gramlich T, Ong JP, Hurley M, *et al.* The utility of radiological imaging in nonalcoholic fatty liver disease. *Gastroenterology* 2002;123:745-50.
9. Lerner RM, Parker KJ, Holen J, Gramiak R, Waag RC. Sono-elasticity: Medical elasticity images derived from ultrasound signals in mechanically vibrated targets. In: *Acoustical Imaging*. Vol. 16. New York: Plenum Press; 1988. p. 317-27.
10. Parker KJ, Huang SR, Musulin RA, Lerner RM. Tissue response to mechanical vibrations for son elasticity imaging. *Ultrasound Med Biol* 1990;16:241-6.
11. Ophir J, Cespedes I, Ponnekanti H, Yazdi Y, Li X. Elastography: A quantitative method for imaging the elasticity of biological tissues. *Ultrasound Imaging* 1991;13:111-34.
12. Lerner RM, Huang SR, Parker KJ. Sonoelasticity images derived from ultrasound signals in mechanically vibrated tissues. *Ultrasound Med Biol* 1990;16:231-9.
13. Sarvazyan AP, Rudenko OV, Swanson SD, Fowlkes JB, Emelianov SY, *et al.* Shear wave elasticity imaging: A new ultrasonic technology of medical diagnostics. *Ultrasound Med Biol* 1998;24:1419-35.
14. Bercoff J, Tanter M, Fink M. Supersonic shear imaging: A new technique for soft tissue elasticity mapping. *IEEE Trans Ultrason Ferroelectr Freq Control* 2004;51:396-409.
15. Trahey GE, Palmeri ML, Bentley RC, Nightingale KR. Acoustic radiation force impulse imaging of the mechanical properties of arteries: *In vivo* and *ex vivo* results. *Ultrasound Med Biol* 2004;30:1163-71.
16. Catheline S, Thomas JL, Wu F, Fink MA. Diffraction field of a low frequency vibrator in soft tissues using transient elastography. *IEEE Trans Ultrason Ferroelectr Freq Control* 1999;46:1013-9.
17. Sandrin L, Tanter M, Gennisson JL, Catheline S, Fink M. Shear elasticity probe for soft tissues with 1-D transient elastography. *IEEE Trans Ultrason Ferroelectr Freq Control* 2002;49:436-46.
18. Catheline S, Gennisson JL, Delon G, Fink M, Sinkus R, Abouelkaram S, *et al.* Measurement of viscoelastic properties of homogeneous soft solid using transient elastography: An inverse problem approach. *J Acoust Soc Am* 2004;116:3734-41.
19. Sandrin L, Tanter M, Catheline S, Fink M. Shear modulus imaging with 2-D transient elastography. *IEEE Trans Ultrason Ferroelectr Freq Control* 2002;49:426-35.
20. Sandrin L, Fourquet B, Hasquenoph JM, Yon S, Fournier C, Mal F, *et al.*

- Transient elastography: A new noninvasive method for assessment of hepatic fibrosis. *Ultrasound Med Biol* 2003;29:1705-13.
21. Nightingale K, McAleavey S, Trahey G. Shear-wave generation using acoustic radiation force: *In vivo* and *ex vivo* results. *Ultrasound Med Biol* 2003;29:1715-23.
  22. Yin M, Talwalkar JA, Glaser KJ, Manduca A, Grimm RC, Rossman PJ, *et al.* Assessment of hepatic fibrosis with magnetic resonance elastography. *Clin Gastroenterol Hepatol* 2007;5:1207-13.e2.
  23. Chen J, Talwalkar JA, Yin M, Glaser KJ, Sanderson SO, Ehman RL. Early detection of nonalcoholic steatohepatitis in patients with nonalcoholic fatty liver disease by using MR Elastography. *Radiology* 2011;259:749-56.